

Effects of droplet fluctuations on the scattering of neutrons and light by microemulsions

V. Lisý and B. Brutovský

Department of Biophysics, P. J. Šafárik University

Jesenná 5, 04154 Košice, Slovakia

e-mail: lisy@kosice.upjs.sk

February 1, 2008

Abstract

Beginning from the first neutron spin-echo study of the shape fluctuations of microemulsion droplets [J. S. Huang, S. T. Milner, B. Farago, and D. Richter, Phys. Rev. Lett. **59**, 2600 (1987)] these experiments are incorrectly interpreted in the literature. This is due to an inappropriate account for the fluctuations and the erroneous application of the original theory to the description of the experiments (see [V. Lisý and B. Brutovsky, Czech J. Phys. **50**, 239 (2000)]). In the presented work both these shortcomings are corrected. We develop the theory of static and dynamic light and neutron scattering from droplet microemulsions. The fluids inside and out of the droplets are separated by a surfactant layer of arbitrary thickness. The scattering functions consistently take into account thermal fluctuations of the shapes of such double-layered spheres to the second order in the changes of their radius. The relaxation times and correlation functions of the fluctuations are found within the Helfrich's theory of interfacial elasticity. The theory is applied to the quantitative description of small-angle neutron scattering, neutron spin-echo spectroscopy and dynamic light scattering experiments. Basic characteristics of the microemulsions, extracted from the fits to the experimental data, significantly differ from those determined in the original works. We include into the consideration the viscosity of the surface layer and give its estimation for the octane - $C_{10}E_5$ - water microemulsion.

Introduction

Microemulsions, thermodynamically stable homogeneous dispersions of oil and water, are intensively studied because of their interesting physical properties and practical importance [1]. The properties of the microemulsions are to a large degree determined by the characteristics of the interface between the bulk fluids. Within the commonly accepted Helfrich's phenomenology [2], this surface layer is described by a few basic parameters. For a microemulsion droplet these are the bending and Gaussian moduli κ and $\bar{\kappa}$, the spontaneous curvature

C_s , the surface tension coefficient α , and the equilibrium radius of the droplet, R_0 . The determination of these characteristics has been attempted by several methods [3]. However, different approaches yield very different values of the parameters of the surface film. So, the neutron spin-echo (NSE) technique of the quasielastic neutron scattering gives significantly larger values κ than indirect macroscopical or optical techniques [3]. This coefficient is in the NSE experiments obtained from the analysis of the peak observed in the Q (wave-vector transfer at the scattering) dependence of the effective diffusion coefficient of the droplets, $D_{eff}(Q)$. The characteristics of the peak are connected with the thermal shape fluctuations of the droplets [3, 4]. In the present work the role of these fluctuations is carefully studied and it is shown that their influence on the observed quantities was underestimated in the interpretation of the scattering experiments. We calculate the intermediate scattering function $F(Q, t)$ that is directly measured by NSE and suitable for the description of small-angle neutron scattering (SANS) and dynamic scattering of light (DLS) as well. As distinct from the previous works, $F(Q, t)$ is found taking into account all the contributions up to the second order of the fluctuations of the droplet radius.

The scattering functions

The microemulsion droplet is modeled by a double-layered sphere with the surface layer of a constant thickness $d = R_2 - R_1$, where $R_1 = R_0 - d/2$ is its inner and R_2 outer radius. Inelastic scattering of neutrons is determined by the time evolution of the scattering length density $\rho(\mathbf{r}, t)$ [5] (in the case of DLS, $\rho(\mathbf{r}, t)$ is replaced by the dielectric constant $\epsilon(\mathbf{r}, t)$ [6]). It is assumed that the scattering from individual droplets is essential. Let the scattering length density is ρ_1 in the droplet interior, ρ_2 in its exterior, and the surface shell is characterized by the constant ρ_0 . The deviation from the radius R_0 of the nondeformed shell (which is also, within the approximation used below, the equivalent-volume radius) is described by the quantity $u = R(t) - R_0$. The quasielastic scattering of neutrons in Born approximation is given by the Van Hove function for coherent scattering, $F(Q, t) = \langle \rho(\mathbf{Q}, t) \rho^*(\mathbf{Q}, 0) \rangle$. To calculate this correlator, u/R_0 is expanded in spherical harmonics [7], with the coefficients $u_{lm}(t)$, where $m = -l, -l+1, \dots, l$, and $l = 0, 1, \dots, l_{max} \sim R_0/a$ (a is a typical molecular diameter). The fluctuations of the molecules in the layer are neglected since the layer is usually thought to be almost incompressible [7]. The shape fluctuations with different numbers l and m are uncorrelated and $\langle u_{lm} \rangle = 0$ for $l > 1$. Using the known properties of the spherical harmonics and spherical Bessel functions j_l , we obtain to the second order in the fluctuations ($Q \neq 0$)

$$\frac{F(Q, t)}{(4\pi R_0^2 d \Delta)^2} = \Phi^2 + 2\Phi\Phi_0 \frac{\langle u_{00} \rangle}{\sqrt{4\pi}} + \sum_{l>1} \frac{2l+1}{4\pi} \left\{ \Phi\Psi \langle u_{l0}^2 \rangle + \Phi_l^2 \langle u_{l0}(0) u_{l0}(t) \rangle \right\}, \quad (1)$$

$$\Phi(Q) = \frac{\rho(Q)}{4\pi R_0^2 d \Delta}, \quad \rho(Q) = \int e^{i\mathbf{Q}\cdot\mathbf{r}} \rho(\mathbf{r}) d\mathbf{r} = 4\pi \left\{ (\rho_1 - \rho_0) R_1^3 \frac{j_1(x_1)}{x_1} + (\rho_0 - \rho_2) R_2^3 \frac{j_1(x_2)}{x_2} \right\},$$

$$\Phi_l(Q) = \frac{1}{R_0 d \Delta} \left[(\rho_1 - \rho_0) R_1^2 j_l(x_1) + (\rho_0 - \rho_2) R_2^2 j_l(x_2) \right],$$

$$\Psi(Q) = \frac{1}{d\Delta} [(\rho_1 - \rho_0)R_1\varphi(x_1) + (\rho_0 - \rho_2)R_2\varphi(x_2)], \quad \varphi(x) = 2j_0(x) - xj_1(x),$$

where $x_{1,2} = QR_{1,2}$, and $\Delta = (\rho_1 + \rho_2)/2 - \rho_0$ characterizes the contrast between the densities of the bulk fluids and the surface shell. The $l = 1$ mode automatically dropped out during the calculations. This coincides with the fact known for the droplets with thin adsorbed layers [7] that this mode corresponds to their translational motion. For $l = 1$ there is no motion in the layer and the droplet moves as a hard sphere [8, 9]. The calculation was carried out in the system connected with the droplet; as usually, the effect of translational motion is included assuming the statistical independence of the translational and other degrees of freedom. Then the right-hand side of eq. (1) has to be multiplied by the factor $\exp(-Q^2Dt)$, where D is the self-diffusion coefficient of a sphere with the hydrodynamic radius $R_H = R_2 + \delta$ (δ accounts for the shell of solvent molecules that move together with the droplet).

Equation (1) is the basic result of the paper. The first term in its right side determines the formfactor of the scattering in the absence of fluctuations [10]. The next terms are connected with the shape fluctuations. For the first time all the second-order terms in the fluctuating density (or dielectric constant) expansion are taken into account. Due to this eq. (1) significantly differs from the expressions for the function $F(Q, t)$ known from the literature. The main difference is in the second term that is given by the $l = 0$ mode contribution. This term did not appear in previous works, though for incompressible bulk fluids $\langle u_{00} \rangle = -\Sigma(2l + 1)\langle u_{l0}u_{l0} \rangle / \sqrt{4\pi}$ ($l > 1$) is nonzero [7]. Also the first term in the sum in eq. (1) usually absents in the description of the scattering experiments [3]. The correct expression for this contribution was found by Gradzielski et al. [11] for the conditions of the perfect shell contrast (deuterated bulk fluids with the densities $\rho_1 \approx \rho_2$ significantly different from the density of the hydrogenated layer). In all the works the scattering function was calculated for d small in comparison with the droplet radius and does not contain the terms $\sim (\rho_1 - \rho_2)$. It is thus inappropriate for the description of the scattering in conditions far from the shell contrast. In the $d/R_0 \rightarrow 0$ limit the previous expressions for $F(Q, t)$ converge to 0 so they cannot describe the scattering from large emulsion or vesicle droplets. The last term in eq. (1) agrees with that from the literature for small d/R_0 and $\rho_1 = \rho_2$.

Equation (1) must be completed by expressions for the correlation functions of the fluctuating droplet radius. In the harmonic approximation for isothermal fluctuations, small thicknesses of the surface layer, and the conservation of the droplet volume as well as the total number of molecules in the shell, we have [12]

$$\langle u_{l0}(t)u_{l0}(0) \rangle \approx \frac{k_B T}{\alpha_l R_0^2(l+2)(l-1)} \exp(-\omega_l t), \quad \alpha_l = \alpha - 2\kappa C_s R_0^{-1} + \kappa l(l+1)R_0^{-1}. \quad (2)$$

The decay rates of the fluctuations, ω_l , have been studied in a number of papers [8, 9]. According to Fujitani [13], the relaxation modes of incompressible dissipative layers are described by the formula

$$\omega_l = \frac{\alpha_l}{R_0} \frac{l(l+1)(l-1)(l+2)}{4(l-1)(l+2)\eta_0/R_0 + (l-1)(2l^2+5l+5)\eta_1 + (l+2)(2l^2-l+2)\eta_2}, \quad (3)$$

where $\eta_{1,2}$ are the viscosity coefficients of the bulk fluids and η_0 is the viscosity of the layer. Finally, the experimental samples contain droplets with different radii so that the measured quantities are the averages over the distribution of the droplets in radii. Within the phenomenological theory of the droplet formation, the distribution function $f(R_0)$ has a form $f(R_0) \sim \exp[-(1 - R_0/R_m)^2/2\epsilon]$, where R_m is the mean radius of the droplets, $\langle (R_0 - R_m)^2 \rangle \approx \epsilon R_m^2$, and ϵ is the polydispersity functionally related to the characteristics of the layer [12]. In the simplest case of the so-called two-phase coexistence $\epsilon = k_B T / 8\pi(2\kappa + \bar{\kappa})$. Note that the entropy-of-mixing contribution is easily incorporated in this relation by adding $2k_B T F(\phi)$ to the denominator [11]. In the random mixing approximation, for small droplet volume fractions ϕ , the function $F(\phi) \approx \ln \phi - 1$. Analogously, eqs. (2) and (3) can be appropriately changed by replacing $R_0^2 \alpha_l \rightarrow \kappa(l-1)(l+2) - \bar{\kappa} - k_B T F(\phi)/4\pi$. This allows us to express the mean quadrate of the fluctuations in the form

$$\langle u_{l0}^2 \rangle = \left\{ (l-1)(l+2) \left[\frac{\kappa}{k_B T} l(l+1) - \frac{1}{8\pi\epsilon} \right] \right\}^{-1} \quad (4)$$

that is particularly suitable to be used in fits to experimental data since it is invariant with respect to the function $F(\phi)$.

The analysis of eqs. (1 - 3) shows that for a number of microemulsion systems studied in the literature the effect of the fluctuations on the scattering functions is important. The fluctuations can significantly affect the position of the minimum in the Q dependence of the scattering intensity measured by SANS, the position of the peak in $D_{eff}(Q)$ observed by NSE, as well as the diffusion coefficient determined using DLS. As a result some basic characteristics of the microemulsions, extracted by us from these experiments, notably differ from those found in the works where the fluctuations were ignored (SANS) [3, 11] or where the account for them should be corrected (NSE, DLS) [3, 4].

Quantitative description of the scattering experiments

We applied the obtained formulae to describe several light and neutron scattering experiments. In the case of large emulsion droplets (the limit $d \rightarrow 0$) eq. (1) well describes the light diffusing-wave spectroscopy experiments [14, 15]. Here we present the results of the description of the experiments on a microemulsion system [3] that well illustrate the importance of the correct account for the effects of fluctuations. In the work [3] the octane- d_{18} - surfactant ($C_{10}E_5$) - water microemulsion in the conditions of two-phase coexistence was studied using SANS, NSE and DLS. The SANS experiments were interpreted using the model of nonfluctuating double-layered spheres [10]. For a diluted microemulsion dispersion the interaction between the droplets was negligible in the region of $Q > 0.03 \text{ \AA}^{-1}$ so that the static structure factor $S(Q) \approx 1$. We have thus calculated the SANS intensity according to the formula $I(Q) \approx$

$N \langle F(Q, 0) \rangle$ [3], where N is the number density of the droplets and the average is over the distribution $f(R_0)$. Figure 1 shows the intensity $I(Q)$ calculated in the case of nonfluctuating droplets and for fluctuating droplets characterized by the parameters determined in [3]. It is seen that both the curves do not describe the experimental data. From our fit to the experiment using the scattering length densities $\rho_1 = 6.35 \times 10^{10}$, $\rho_2 = 6.36 \times 10^{10}$ [11] and $\rho_0 = 1.65 \times 10^9 \text{cm}^{-2}$ [16], we obtained the set of the parameters R_m , d , ϵ and κ for which the theoretical curve well reproduces the measured $I(Q)$.




Figure 1: The SANS intensity calculated from eqs. (1-2) with (the upper full line) and without (dashed line) the droplet fluctuations in the shape. The experimental points are from the work [3]. The octane- d_18 - surfactant ($C_{10}E_5$) - water microemulsion at the temperature $T = 305.2$ K is characterized by the parameters $\rho_1 = 6.35 \times 10^{10}$, $\rho_2 = 6.36 \times 10^{10}$ and $\rho_0 = 1.65 \times 10^9 \text{cm}^{-2}$, $\kappa = 0.92 k_B T$, $\epsilon = 0.036$, $d = 10.8 \text{\AA}$, $R_m = 48 \text{\AA}$, and the volume fraction of the droplets is 0.049 [3]. Our fit to the experimental data yielded the values $\kappa = 1.93 k_B T$, $\epsilon = 0.046$, $R_m = 47.7 \text{\AA}$ and $d = 6.4 \text{\AA}$.

With these parameters and for dielectric constants $\epsilon_1 = 1.946$, $\epsilon_2 = 1.769$ and $\epsilon_0 = 2.106$ [16], we evaluated the effective diffusion coefficient of the droplets, $D_{eff} = - \langle dF(Q, t)/dt \rangle / (Q^2 \langle F(Q, 0) \rangle)$ at $t = 0$. By DLS this coefficient is measured in the region of Q much smaller than in the case of SANS. In this region D_{eff} is practically constant, not influenced by the time-dependent terms $\sim \langle u(0)u(t) \rangle$ in eq. (1). The Stokes self-diffusion coefficient has a form $D = k_B T / 6\pi\eta_2 R_H$, $R_H = R_0 + d/2 + \delta$, and its mean value was determined from the requirement to satisfy the measured $D_{eff} = 3.8 \times 10^{-11} \text{m}^2/\text{s}$ (with the tolerance $\pm 5\%$). Our attempt to describe the DLS experiment assuming a hydration shell with a small δ (2-3 $\text{\AA}$) was not possible without a change of the parameters R_m , d , ϵ and κ , that makes worse the agreement with the SANS experiment. The best agreement with both the experiments required $\delta = 13 \div 21 \text{\AA}$ (the mean $\delta \approx 17 \text{\AA}$). The found thickness of the hydration shell assumes

the shell built of several layers [3]. Such a possibility is a question for future investigations. We suppose however that first a model taking into account different properties of the bound and free bulk fluids should be developed. Finally, we turned to the NSE experiment. This experiment was in the work [3] described using the scattering function (1) without the time-independent contributions of the fluctuations and for the ideal contrast. The decay rates were from our paper [17], however, those ω_l correspond to infinitely easily compressible layers [8, 9]. In our analysis we used ω_l from eq. (3) appropriate for incompressible layers. The viscosities of the bulk fluids were $\eta_1 = 0.4675$ and $\eta_2 = 0.7632$ cP at $T = 305.9$ K [3]. The last parameter needed to describe the experiment is η_0 . If the previous theories are correctly used the height of the peak in D_{eff} is several times larger than its measured values. Also in our theory this height is too large. We propose to solve this problem introducing the dissipation in the layer that leads to a decrease of ω_l and thus to a decrease of the peak height. From the correspondence between the experimental [3] and theoretical heights of the peak we found $\eta_0 = 5.6 \times 10^{-11}$ Ns/m, the value only weakly influenced by the parameter δ . The calculated and experimental D_{eff} is shown on Figure 2. The rest of the parameters estimated from the fits are as follows: $\kappa = 1.93(0.92)k_B T$, $\epsilon = 0.046(0.036)$, $\bar{\kappa} = -2.68(-0.38)k_B T$, $R_m = 47.7(48)\text{\AA}$ and $d = 6.4(10.8)\text{\AA}$ (in parentheses the values from [3] are shown). When the entropy-of-mixing contributions are taken into account, only the parameter $\bar{\kappa}$ slightly changes, now being $\bar{\kappa} = -3k_B T$. The hydrodynamic radius of the droplets and the diffusion coefficient are about $R_H = 68(81)\text{\AA}$ and $D(R_m) = 4.3(3.6) \times 10^{-11} \text{m}^2/\text{s}$.




Figure 2: The effective diffusion coefficient measured by NSE (points) [3] on the same system as in Fig. 1, and calculated from eqs. (1-4) using the parameters obtained from the fit to SANS (Fig. 1). The height of the peak corresponds to the experimental one if the viscosity of the surface layer $\eta_0 = 5.6 \times 10^{-11} \text{Ns/m}$ is employed in the calculations. The thickness of the hydration shell $\delta = 17\text{\AA}$ was obtained from the DLS experiment [3]. The viscosities of the bulk fluids were $\eta_1 = 0.4675$ and $\eta_2 = 0.7632$ cP.

The difference between the obtained parameters and those found in the original work [3] concerns first of all the elastic coefficients that are the crucial parameters needed to describe the properties of microemulsions. For the first time we have from the scattering experiments estimated the viscosity coefficient of the surface layer.

In conclusion, a deeper consideration of the effects of the shape fluctuations of microemulsion droplets on the static and dynamic neutron and light scattering was presented. The agreement with such experiments is much better than it was possible so far. More reliable information can be now obtained about important phenomenological parameters of microemulsions. We suppose that in further investigations the role of the hydration layer bound to the droplets should be studied in detail. More should be also known about the interaction between the droplets and the entropy of dispersion that are differently treated in the literature [11, 18].

This work was supported by the grant No. 1/7401/20, VEGA, Slovak Republic.

References

- [1] D. O. Shah, editor. *Micelles, Microemulsions, and Monolayers*, M. Dekker, New York, 1998
- [2] W. Helfrich, *Z. Naturforsch.* **28c**, 693 (1973)
- [3] T. Hellweg and D. Langevin, *Phys. Rev. E* **57**, 6825 (1998)
- [4] B. Farago, *Physica B* **51**, 226 (1996)
- [5] S. W. Lovesey and P. Schofieldm, *J. Phys. C* **9**, 2843 (1976)
- [6] L. I. Komarov and I. Z. Fisher, *Zh. Eksp. Theor. Fiz.* **43**, 1927 (1962)
- [7] S. T. Milner and S. A. Safran, *Phys. Rev. A* **36**, 4371 (1987)
- [8] V. Lisý, B. Brutovský, and A. V. Zatovsky, *Phys. Rev. E* **58**, 7598 (1998)
- [9] V. Lisý, B. Brutovský, P. Miškovský, and A. V. Zatovsky, *Czech. J. Phys.* **49**, 137 (1999)
- [10] A. Guinier and G. Fournet, *Small-Angle Scattering of X-Rays*. Wiley, New York, 1955
- [11] M. Gradzielski, D. Langevin, and B. Farago, *Phys. Rev. E* **53**, 3900 (1996)
- [12] V. Lisý and B. Brutovský, *Czech. J. Phys.* **50**, 239 (2000)
- [13] Y. Fujitani, *Physica A* **203**, 214 (1994)
- [14] V. Lisý and B. Brutovský, *Phys. Rev. E* **59**, 3765 (1999)
- [15] V. Lisý, B. Brutovský, and P. Miškovský, *Physica B* **266**, 300 (1999)
- [16] D. Langevin and Lay-Theng Lee, personal communications
- [17] V. Lisý, *Phys. Lett. A*, **150** 105 (1990)
- [18] E. Ruckenstein, *J. Coll. Interface Sci.* **204**, 143 (1998)